

Extraterrestrial chromite in latest Maastrichtian and Paleocene pelagic limestone at Gubbio, Italy: The flux of unmelted ordinary chondrites

Anders CRONHOLM* and Birger SCHMITZ

Department of Geology, Lund University, Sölvegatan 12, SE-22362 Lund, Sweden

*Corresponding author. E-mail: anders.cronholm@geol.lu.se

(Received 17 January 2007; revision accepted 20 June 2007)

Abstract—The distribution of sediment-dispersed extraterrestrial (ordinary chondritic) chromite (EC) grains ($>63\ \mu\text{m}$) has been studied across the latest Maastrichtian and Paleocene in the Bottaccione Gorge section at Gubbio, Italy. This section is ideal for determining the accumulation rate of EC because of its condensed nature and well-constrained sedimentation rates. In a total of 210 kg of limestone representing eight samples of 14–28 kg distributed across 24 m of the Bottaccione section, only 6 EC grains were found (an average of $0.03\ \text{EC grains kg}^{-1}$). In addition, one probable pallasitic chromite grain was found. No EC grains could be found in two samples at the Cretaceous-Tertiary (K-T) boundary, which is consistent with the K-T boundary impactor being a carbonaceous chondrite or comet low in chromite. The average influx of EC to Earth is calculated to $\sim 0.26\ \text{grain m}^{-2}\ \text{kyr}^{-1}$. This corresponds to a total flux of ~ 200 tons of extraterrestrial matter per year, compared to $\sim 30,000$ tons per year, as estimated from Os isotopes in deep-sea sediments. The difference is explained by the EC grains representing only unmelted ordinary chondritic matter, predominantly in the size range from $\sim 0.1\ \text{mm}$ to a few centimeters in diameter. Sedimentary EC grains can thus give important information on the extent to which micrometeorites and small meteorites survive the passage through the atmosphere. The average of $0.03\ \text{EC grain kg}^{-1}$ in the Gubbio limestone contrasts with the up to $\sim 3\ \text{EC grains kg}^{-1}$ in mid-Ordovician limestone that formed after the disruption of the L chondrite parent body in the asteroid belt at $\sim 470\ \text{Ma}$. The two types of limestone were deposited at about the same rate, and the difference in EC abundance gives support for an increase by two orders of magnitude in the flux of chondritic matter directly after the asteroid breakup.

INTRODUCTION

Recent empirical evidence indicates the existence of short periods during the Phanerozoic with significantly increased flux of extraterrestrial matter to Earth (Farley et al. 1998, 2006; Schmitz et al. 1996, 2001, 2003). Such flux increases have been predicted to result from, for example, gravity perturbations of the orbits of comets in the Oort cloud (e.g., Hills 1981; Hut et al. 1987; Matese et al. 1995) or parent body breakup events in the asteroid belt (Zappalà et al. 1998; Nesvorný et al. 2006). The finding of more than fifty fossil meteorites in mid-Ordovician marine limestone at Kinnekulle in southern Sweden supports an increase in the flux of meteorites by one to two orders of magnitude following the disruption of the L chondrite parent body in the asteroid belt $\sim 470\ \text{Myr}$ ago (Schmitz et al. 1996, 2001). The fossil meteorites have been shown to be pseudomorphosed L chondrites (Schmitz et al. 2001; Bridges et al. 2007; Greenwood et al. 2007) and their relict chromite grains show

cosmic-ray exposure ages that support an origin from an asteroid breakup event (Heck et al. 2004). A mid-Ordovician increase in the meteorite flux is further supported by the distribution of sediment-dispersed extraterrestrial (ordinary chondritic) chromite (EC) grains ($63\text{--}355\ \mu\text{m}$) in condensed limestone strata from southern Sweden (Schmitz et al. 2003; Schmitz and Häggström 2006). The search for fossil meteorites is time-consuming and requires screening of thousands of cubic meters of limestone, whereas sediment-dispersed EC grains can be searched in samples of 10–50 kg. Chromite is highly resistant to weathering and it is one of the rare relict minerals of fossil meteorites (Thorslund et al. 1984). The sediment-dispersed EC grains represent residues of primarily small, $0.1\ \text{mm}$ to $1\text{--}3\ \text{cm}$ ordinary chondrites that decomposed on the sea floor (Schmitz et al. 2003; Heck et al., Forthcoming). Recent ordinary chondrites contain on average $\sim 0.25\ \text{wt\%}$ chromite (FeCr_2O_4) (Keil 1962) and constitute $\sim 80\%$ of the total meteoritic influx (Bevan et al. 1998). The EC can be readily distinguished from terrestrial chromite by a




Fig. 1. Geology of Gubbio area and location of Bottaccione Gorge section and outcrop studied. Modified after Alvarez et al. (1977).

characteristic and unique major and trace element composition (Thorslund et al. 1984; Nyström et al. 1988; Roeder 1994; Schmitz et al. 2001). Extraterrestrial chromite shows minor variations in composition related to types H, L, and LL of host meteorite (Bunch et al. 1967; Schmitz et al. 2001; Wlotzka 2005). Hence, EC grains dispersed in sediments can be useful for determining both ancient accretion rates of ordinary chondritic material and variations in the types of ordinary chondrites reaching Earth (Schmitz et al. 2003).

In mid-Ordovician condensed strata in southern Sweden, limestone beds that formed after the disruption of the L chondrite parent body contain typically 1–3 EC grains per kg limestone, whereas in 379 kg of limestone that formed during the ~5 Myr immediately prior to this event, only 5 EC grains were found (Schmitz and Häggström 2006). In order to fully assess the significance of these results, EC accretion rates also for other periods in Earth's history must be determined. In the present study the classic section across the latest Maastrichtian and Paleocene in the Bottaccione Gorge at Gubbio, Italy, was selected for EC searches. The pelagic limestone at this site formed at the same low sedimentation rates, i.e., a few millimeters per 1000 years, as the mid-Ordovician limestone at Kinnekulle (Arthur and Fischer

1977; Schmitz et al. 1996). Detailed bio- and magnetostratigraphies have been established for the Bottaccione Gorge section showing that it is essentially complete over the stratigraphic interval considered here (e.g., Alvarez et al. 1977; Premoli Silva 1977; Monechi and Thierstein 1985; Cresta et al. 1989; Premoli Silva and Sliter 1995), thus making it possible to constrain average sedimentation rates with great confidence. A detailed study of ^3He across the section has determined the variations in the flux of interplanetary dust particles to Earth (Mukhopadhyay et al. 2001a, 2001b). Helium-3 records primarily reflect the influx of the smallest extraterrestrial particles, in the size range of ~3–35 μm , with an origin from both comets and asteroids.

The Gubbio section is thus ideal for establishing accretion rates for EC; additionally, light can be shed on whether the Cretaceous-Tertiary (K-T) boundary asteroid or comet impact (Alvarez et al. 1980) was associated with any increase in the flux of ordinary chondritic matter to Earth. It should be noted that our approach differs from previous studies of cosmic spinels at the K-T boundary, mainly since these studies focused on spinels (~1–50 μm) formed during impacts of cosmic objects and/or atmospheric entry of cosmic materials (Smit and Kyte 1984; Kyte and Smit 1986; Robin et al. 1992; Toppani and Libourel 2003; Ebel and Grossman 2005) rather than the coarse (>63 μm) and common chromite originally present in equilibrated ordinary chondrites (Keil 1962; Ramdohr 1973; Rubin 1997; Bridges et al. 2007). In unequilibrated ordinary chondrites, i.e., petrological type 3, the chromite grains are usually smaller than 63 μm (Bridges et al. 2007) and hence lie outside the range of detection in the present study.

GEOLOGICAL SETTING

The Bottaccione Gorge with its famous and well-studied pelagic sequence of the middle Cretaceous to the Eocene (e.g., Arthur and Fischer 1977; Alvarez et al. 1980) is located in the Umbria-Marche Apennines, central Italy, directly north of the medieval town of Gubbio (43°22'N, 12°35'E, along the state road SS298 to Scheggia; see Fig. 1). Our sampled interval belongs to the ~325 m thick Scaglia Rossa Formation, which extends from the early Turonian to the middle Eocene (Arthur and Fischer 1977; Monechi and Thierstein 1985). The upper part of this formation consists predominantly of pink to red homogenous limestone and marly limestone that have been thoroughly bioturbated and compacted prior to cementation. The pristine condition of the Scaglia Rossa is explained by the apparent absence of intrusions and thermal alterations in the area (Arthur and Fischer 1977). Agglutinated foraminifera indicate that during the late Cretaceous and Paleocene, the Gubbio/Umbria-Marche basin had a deep bathyal position with a water depth of 1500–2500 m (Kuhnt 1990). Pelagic conditions are also




Fig. 2. Distribution of samples studied and number of recovered extraterrestrial (EC) and other (OC) chromite grains across the Bottaccione Gorge section. Lithology after Arthur and Fischer (1977), helium isotopic distribution from Mukhopadhyay et al. (2001), foraminifera stratigraphy from Premoli Silva (1977) and Premoli Silva and Sliter (1995), and nannofossil stratigraphy from Monechi and Thiersten (1985).

indicated by abundant calcareous nannofossils and foraminifera, an aeolian clay and silt component, high planktic/benthic ratio in foraminifera, and extremely low concentrations of organic carbon (Arthur and Fischer 1977; Montanari 1991). In about 50 samples spanning from the Turonian to early Paleocene, Kuhnt (1990) noted the absence of terrigenous detrital material in grain fractions larger than 63 μm .

It is generally considered that the sedimentation rates were extremely low in the Gubbio area from the top Cretaceous (6–10 mm kyr^{-1}) to the lowermost Paleocene limestone (2–5 mm kyr^{-1}) (Montanari and Koeberl 2000). For the Paleocene epoch, Arthur and Fischer (1977) suggest an average sedimentation rate of 2.4 mm kyr^{-1} . Mukhopadhyay et al. (2001b) suggest average sedimentation rates on the order of 4–6 mm kyr^{-1} for the latest Cretaceous and 2–

3 mm kyr^{-1} for the early and middle Paleocene. Their helium isotopic data show that the flux of extraterrestrial ^3He to Earth is constant within a factor of ~ 2 across the K-T boundary and the early and middle Paleocene. Bulk rock ^3He concentrations vary by a factor of ~ 4 , reflecting mainly short-term variations in sedimentation rates (Fig. 2).

The Bottaccione Gorge section became famous as the first section shown to contain enhanced iridium levels (3–6 ppb) at the K-T boundary (Alvarez et al. 1980). The iridium originates from a carbonaceous chondritic asteroid or a comet (diameter of ~ 10 km) that impacted on the Yucatán Peninsula, Mexico (Hildebrand et al. 1991; Shukolyukov and Lugmair 1998). The K-T boundary in the Bottaccione Gorge section is marked by a 1–2 cm thick dark clay layer underlain by a white bleached limestone zone, ~ 20 –50 cm thick. This zone is related to more reducing sediment conditions when increased

amounts of organic material reached the sea floor during the K-T boundary event (Lowrie et al. 1990; Montanari and Koeberl 2000). With the exception of the thin foraminiferal P0 zone, the Bottaccione Gorge section contains an essentially complete succession of biozones across the K-T transition (Smit 1982). The P0 zone is absent or unrecognizable in most of the Umbria-Marche locations, possibly reflecting slow sedimentation rates combined with thorough bioturbation, which would mix tests from the thin P0 zone and the overlying foraminiferal zones (Montanari and Koeberl 2000). Previous studies of the K-T boundary clay worldwide have located excess amounts of impact-related cosmic spinels and shocked quartz, but according to Montanari and Koeberl (2000), “despite its fame . . . the Bottaccione K-T clay is one of the worst known in the Umbria-Marche region in terms of preservation and suitability for a high-resolution microstratigraphic study.” The clay is affected by shearing caused by flexural slip folding and suffers from contamination by the surrounding soil. Common spinel-bearing spherules and shocked quartz have instead been found in the best preserved boundary clay at Petriccio only ~30 km north of Gubbio (Montanari 1991). However, this site is no longer easily accessible.

MATERIALS AND METHODS

Seven large limestone samples ranging from 27.7 to 28.3 kg plus one sample of 13.8 kg, for a total of 210 kg, were collected from 7.15 m below the K-T boundary clay to 17.18 m above it (Fig. 2). One sample of 28.3 kg spanned the 20 cm of section starting 1 cm above the base of the K-T boundary clay; another sample of 13.8 kg spanned the 10 cm of bleached strata immediately underlying the clay. The remaining samples (depths in meters relative to the K-T boundary clay: -7.15 to -6.90, -0.90 to -0.65, +2.10 to +2.40, +5.98 to +6.05, +11.15 to +11.40, +16.85 to +17.18 m) were regarded as representing “background levels” not influenced by the K-T boundary event. The stratigraphically lowest sample is dated to the planktonic foraminifera *Abathomphalus mayaroensis* zone of the latest Maastrichtian (Premoli Silva 1977). The youngest sample at 17 m is dated to the lowest part of the planktonic foraminiferal P4 zone (Premoli Silva 1977) or calcareous nannofossil zone NP5 (Monechi and Thierstein 1985) (Fig. 2).

The eight samples studied were decalcified in 6 M hydrochloric acid at room temperature and then sieved at 32 μm . The residual fractions were leached in 18 M hydrofluoric acid at room temperature and separated into three size fractions (32–63, 63–355, and >355 μm) by sieving and drying. Some samples contained common coal particles and were burned at 550 °C for 24 h to destroy the coal. The coal particles are removed since they resemble the chromite grains. Tests have confirmed that the heating does not affect the major and trace element chemistry of the chromite grains.

The middle size fraction (63–355 μm) was scrutinized under a stereomicroscope and suspected chromite grains were collected with a fine brush. The grains were first mounted on carbon tape and preliminarily analyzed in an unpolished state for major and trace elements. The grains were thereafter mounted in epoxy resin and polished before final elemental analysis. For polishing, Struer’s alumina paste (standard quality) mixed with water was used on a spinning polishing cloth (Buehler). For a few grains that were destroyed during the polishing phase, we use the results for unpolished grains. Quantitative elemental analyses were conducted with an energy dispersive spectrometer (INCA x-sight, Oxford Instruments) with a Si detector linked to a Hitachi S-3400N scanning electron microscope (SEM). Samples were analyzed at an acceleration voltage of 15 kV, a beam 1–2 μm in diameter, and a counting live-time of 80 s. Cobalt was used as a standard; the precision (reproducibility) of analyses was typically within 1–4%. Analytical accuracy was controlled by repeated analyses of the USNM 117075 chromite (Smithsonian) reference standard (Jarosewich et al. 1980).

DISTRIBUTION AND ORIGIN OF CHROME SPINELS

The distribution of sediment-dispersed chrome-rich spinel grains (63–355 μm) in the Bottaccione Gorge section is shown in Fig. 2 and the chemical composition is shown in Tables 1–4. We follow the procedures outlined by Schmitz and Häggström (2006) and divide the chrome-rich spinels into two groups: extraterrestrial (equilibrated ordinary chondritic) chromite (EC) and “other” chromium-rich spinels (OC). Under the binocular microscope, the EC and OC grains cannot be distinguished from each other, but they differ in chemical composition. The EC grains are characterized primarily by high Cr_2O_3 contents, ~55–60 wt%, FeO concentrations in the range of ~25–30 wt%, low Al_2O_3 at ~5–8 wt%, and low MgO concentrations of ~1.5–4 wt%. The most discriminative feature, however, is that they plot in narrow ranges of V_2O_3 (~0.6–0.9 wt%) and TiO_2 (~2.0–3.5 wt%) concentration. For a grain to be classified as an EC grain, it has to have a composition within or very close to the defined ranges for all the elements listed above. The OC grains, on the other hand, have a wide compositional range. In most cases they have a terrestrial origin, but some grains may represent rare types of extraterrestrial spinels.

In total, only 6 EC and 14 OC grains were retrieved from 210 kg of limestone (Tables 1–4). No chrome-rich spinels were present in the two samples at the K-T boundary (0 to -0.10 m and +0.01 to +0.20 m). Instead, the 6 EC grains were found in 4 samples evenly distributed through the section. The highest concentrations of EC and OC grains were found in the sample at +2.10 m, which contained 3 EC and 8 OC grains. The average EC and OC concentrations in the section are 0.029 and 0.067 grains kg^{-1} limestone,

Table 1. The element concentration (wt%) of extraterrestrial chromite (EC) grains from this study.

| Gubbio EC grains ^a | Cr ₂ O ₃ | Al ₂ O ₃ | MgO | TiO ₂ | V ₂ O ₃ | FeO | MnO | ZnO | Total |
|-------------------------------|--------------------------------|--------------------------------|-------------|------------------|-------------------------------|--------------|-------------|-------------|-------|
| –0.90 to –0.65 m | #1 | 59.53 | 7.48 | 3.03 | 2.20 | 22.52 | 1.05 | 2.58 | 99.2 |
| +2.10 to 2.40 m | #1 ^b | 56.27 | 5.27 | 6.08 | 1.67 | 28.45 | 0.71 | 0.95 | 100.0 |
| | #2 | 59.91 | 6.30 | 4.78 | 1.86 | 24.92 | n.d. | 1.14 | 99.4 |
| | #3 | 57.38 | 4.30 | 6.94 | 2.36 | 24.14 | 0.89 | 2.24 | 98.9 |
| +5.98 to 6.05 m | #1 | 59.56 | 6.14 | 2.11 | 3.20 | 25.25 | 0.67 | 0.51 | 98.1 |
| +11.15 to 11.40 m | #1 | 60.96 | 5.18 | 3.33 | 2.83 | 23.93 | 0.75 | 0.77 | 98.6 |
| Mean (with s.d.) | | 58.94 ± 1.75 | 5.78 ± 1.11 | 4.38 ± 1.88 | 2.35 ± 0.58 | 24.87 ± 1.99 | 0.68 ± 0.36 | 1.37 ± 0.84 | 99.0 |

^aDepth relative to the K-T boundary in the Bortaccone Gorge section, Gubbio.^bThe grain was lost during polishing, and is only represented by preliminary pre-polish measurements. The elemental values have been normalized from the original total of 92.8 wt%. n.d. = not detected.

Table 2. The average element concentration (wt%, with standard deviation) of EC grains from this and previous studies.

| | Cr ₂ O ₃ | Al ₂ O ₃ | MgO | TiO ₂ | V ₂ O ₃ | FeO | MnO | ZnO | Total |
|--|--------------------------------|--------------------------------|-------------|------------------|-------------------------------|-------------|-------------|-------------|-------|
| Sediment-dispersed EC grains | | | | | | | | | |
| Gubbio section, Italy; 6 grains (this study) | 58.9 ± 1.75 | 5.78 ± 1.11 | 4.38 ± 1.88 | 2.35 ± 0.58 | 0.67 ± 0.12 | 24.9 ± 1.99 | 0.68 ± 0.36 | 1.37 ± 0.84 | 99.0 |
| Thorsberg quarry, Sweden; 276 grains ^a | 57.6 ± 1.58 | 6.07 ± 0.76 | 2.58 ± 0.79 | 3.09 ± 0.33 | 0.75 ± 0.07 | 27.4 ± 2.63 | 0.78 ± 0.20 | 0.53 ± 0.50 | 98.8 |
| Chromite grains from meteorites ^b | | | | | | | | | |
| Mid-Ordovician fossil meteorites, Thorsberg quarry; 26 meteorites | 57.6 ± 1.30 | 5.53 ± 0.29 | 2.57 ± 0.83 | 2.73 ± 0.40 | 0.73 ± 0.03 | 26.9 ± 3.89 | 1.01 ± 0.33 | 1.86 ± 2.43 | 99.0 |
| Recent H5/6 group chondrites; 11 meteorites | 56.6 ± 0.37 | 6.44 ± 0.14 | 2.98 ± 0.23 | 2.20 ± 0.17 | 0.73 ± 0.02 | 29.3 ± 0.67 | 1.00 ± 0.08 | 0.33 ± 0.05 | 99.6 |
| Recent L5/6 group chondrites; 10 meteorites | 56.0 ± 0.65 | 5.97 ± 0.43 | 2.93 ± 0.97 | 2.68 ± 0.40 | 0.75 ± 0.02 | 30.2 ± 2.23 | 0.83 ± 0.10 | 0.30 ± 0.07 | 99.7 |

^aData from Schmitz and Häggström (2006).^bData given is the mean of the average composition of typically 10–20 EC grains from each meteorite. Data from Schmitz et al. (2001), modified by Alwmark and Schmitz (2007).

Table 3. The element concentration (wt%) of other chromium-rich spinel (OC) grains from this study.

| Gubbio OC grains ^a | | Cr ₂ O ₃ | Al ₂ O ₃ | MgO | TiO ₂ | V ₂ O ₃ | FeO | MnO | ZnO | Total |
|-------------------------------|-----------------|--------------------------------|--------------------------------|-------|------------------|-------------------------------|-------|------|------|-------|
| +2.10 to 2.40 m | #1 | 27.61 | 29.54 | 15.47 | 2.08 | n.d. | 23.80 | n.d. | n.d. | 98.5 |
| | #2 | 23.80 | 32.65 | 15.21 | 1.80 | n.d. | 24.19 | n.d. | n.d. | 97.7 |
| | #3 | 26.73 | 31.07 | 16.00 | 1.93 | n.d. | 22.95 | n.d. | n.d. | 98.7 |
| | #4 | 18.06 | 30.21 | 9.70 | 2.50 | n.d. | 36.91 | 0.92 | n.d. | 98.3 |
| | #5 | 11.73 | 38.51 | 14.80 | 3.05 | n.d. | 30.91 | n.d. | n.d. | 99.0 |
| | #6 ^b | 28.67 | 38.67 | 15.96 | 0.26 | n.d. | 16.44 | n.d. | n.d. | 100.0 |
| | #7 ^c | (grain lost during polishing) | | | | | | | | |
| | #8 ^c | (grain lost during polishing) | | | | | | | | |
| +5.98 to 6.05 m | #1 | 64.27 | 5.83 | 5.42 | n.d. | 0.56 | 14.20 | 0.71 | 8.11 | 99.1 |
| +11.15 to 11.40 m | #1 | 43.46 | 27.58 | 12.53 | n.d. | n.d. | 16.34 | n.d. | n.d. | 99.9 |
| | #2 | 26.61 | 27.16 | 12.19 | 1.73 | n.d. | 30.35 | n.d. | n.d. | 98.0 |
| | #3 | 42.13 | 30.74 | 12.74 | n.d. | n.d. | 14.56 | n.d. | n.d. | 100.2 |
| | #4 ^d | (grain lost during polishing) | | | | | | | | |
| +16.85 to 17.18 m | #1 | 56.79 | 15.27 | 8.08 | n.d. | n.d. | 15.00 | n.d. | 2.20 | 97.3 |

^aDepth relative to the K-T boundary in the Bottaccione Gorge section, Gubbio.

^bThe grain was lost during polishing, and is only represented by preliminary pre-polish measurements. The elemental values have been normalized from the original total of 96.4 wt%.

^cPre-polish measurements of grains #7 and 8 gave similar results as for grains #1, 2, and 3 in the same sample.

^dPre-polish measurements of grain #4 gave similar results as for grains #1 and 3 in the same sample.

n.d. = not detected.

Table 4. The element composition (wt%) of average pallasitic chromite compared to the OC grain from +5.98 to 6.05 m above the K-T boundary at Gubbio, Italy.

| | Cr ₂ O ₃ | Al ₂ O ₃ | MgO | TiO ₂ | V ₂ O ₃ | FeO | MnO | ZnO | Total |
|--------------------------------|--------------------------------|--------------------------------|-----|------------------|-------------------------------|------|------|-------|-------|
| Average pallasite ^a | 64.0 | 5.6 | 5.8 | 0.18 | 0.54 | 23.2 | 0.65 | <0.02 | 99.97 |
| OC grain from +5.98 m | 64.3 | 5.8 | 5.4 | n.d. | 0.56 | 14.2 | 0.71 | 8.1 | 99.1 |

^aData from Bunch and Keil (1971).

n.d. = not detected.

respectively. The EC grains from the Gubbio section are on the whole very similar in composition to both chromite in recent ordinary chondrites and EC grains found in Ordovician limestone and fossil meteorites at Kinnekulle, Sweden (Schmitz et al. 2001, 2003; Schmitz and Häggström 2006) (see Table 2). Some minor but significant differences exist. The Gubbio grains are often high in ZnO (0.5–2.6 wt%) and on average 2–2.5 wt% lower in FeO than Ordovician chondritic chromite. For the Ordovician fossil meteorites, it has been shown that Zn can replace Fe in the chromite structure during diagenesis (Schmitz et al. 2001); hence, the Gubbio grains appear to be more diagenetically altered than most Ordovician EC grains. The Gubbio EC grains on average have a higher MgO (4.4 wt%) concentration than the Ordovician (2.6 wt% MgO) EC grains. However, only the three grains from the sample at +2.10 m have significantly higher MgO than the Ordovician average EC. Single fossil meteorites at Kinnekulle contain chromite grains with MgO concentrations of ~5–6 wt% rather than the normal ~2–4 wt% (Schmitz et al. 2001). Hence, the three EC grains from the +2.10 m level probably derive from the same meteorite or micrometeorite. The TiO₂ concentration of the Gubbio EC grains is on average 0.4–

0.7 wt% lower than in Ordovician EC grains. Previous studies have shown that the TiO₂ concentration of sediment-dispersed chromite grains in general is very resistant to weathering and diagenesis (Schmitz et al. 2001; Schmitz and Häggström 2006; Alwmark and Schmitz 2007). Chromite in H chondrites is on average ~0.6–0.7 wt% lower in TiO₂ than in L chondrites (Bunch et al. 1967; Schmitz et al. 2001; Wlotzka 2005). The Ordovician EC grains predominantly originate from the disruption of the L chondrite parent body at this time, which could explain the higher TiO₂. There is no reason to expect that L chondrites dominated the flux in the latest Maastrichtian-Paleocene. Although the majority of the EC grains from Gubbio may derive from H chondrites, the ranges of TiO₂ concentrations in recent H and L chondritic chromite overlap; hence, only the average TiO₂ content of a large number of sediment-dispersed grains is indicative of precursor chondrite type. Moreover, the three grains in the sample at +2.10 m are low in TiO₂ and this affects the average TiO₂ concentration disproportionately. The three EC grains at other levels have typical H or L chondritic TiO₂ contents. In essence, no conclusions about the relative fluxes of H and L chondrites can be drawn from this small data set.

All except 2 of the 14 OC grains originate from the two samples at +2.10 and +11.15 m (Table 3). Most of the OC grains from these samples are similar in composition. They have low Cr/Cr + Al ratios (<0.5), which is in good agreement with a terrestrial origin (Roeder 1994), they are typically high in MgO (12–16 wt%), and they contain no detectable V or Zn. Comparisons of the compositions of the OC grains from Gubbio with chrome spinels from various magmatic provinces as plotted in Barnes and Roeder (2001) indicate an ophiolite provenance, but alternative sources are also plausible. Various magmatic processes, including subduction and possibly plume activity, have occurred in Italy dating back to at least 60–70 Ma (Bell et al. 2004). A more detailed evaluation of the origin of the terrestrial chromite grains from the Gubbio section is beyond the scope of this article.

The two OC grains from the samples at +5.98 and +16.85 m show different compositions compared to those discussed above (Table 3). The origin of the grain from +16.85 m is uncertain and it may be a terrestrial grain. The single OC grain at +5.98 m represents a very special case. It is unusually large ($150 \times 300 \mu\text{m}$) and has a distinct composition almost identical to the average composition of chromites from pallasites (Table 4) (Bunch and Keil 1971). The high Cr_2O_3 (64%), low Al_2O_3 (5.6 wt%), and characteristic V_2O_3 (0.5 wt%) and TiO_2 (0.18 wt%) contents distinguish pallasitic chromite from most terrestrial chromite. The higher ZnO (8.1 wt%) and lower FeO (14.2 wt%) of our OC grain compared to the pallasitic average chromite is readily explained by replacement of Fe by Zn during diagenesis (Schmitz et al. 2001). Pallasitic chromite is also characterized by a larger grain size than in other types of meteorites (Bunch and Keil 1971). The fact that the grain occurs unassociated with any other OC grains in the same sample and the profound difference in composition compared to the terrestrial chrome spinels in the samples above and below also speak in favor of an extraterrestrial origin.

BACKGROUND FLUX OF EXTRATERRESTRIAL CHROMITE

Previous attempts to determine the time-averaged total influx of meteorites and/or smaller extraterrestrial particles to Earth have relied on the density of micrometeoroid hyper-velocity impact craters on the Long Duration Exposure Facility (LDEF) satellite (Love and Brownlee 1993), radar micrometeor observations in the upper atmosphere (Mathews et al. 2001), concentrations of platinum group elements, osmium isotopes and ^3He in condensed sediments (Farley et al. 1997; Peucker-Ehrenbrink and Ravizza 2000; Peucker-Ehrenbrink 2001; Dalai and Ravizza 2006), iridium concentrations in ice cores (Karner et al. 2003; Gabrielli et al. 2004), micrometeorite abundances in Antarctic ice (Taylor and Lever 2001) and, for meteorites, meteor sky-watch programs (Halliday 2001) and systematic meteorite searches

in wind-eroded desert areas (Bland et al. 1996; Bland 2001). The various approaches give flux estimates that sometimes differ significantly. To some extent this can be explained by the different size fractions of the total flux being measured, but methodological problems such as undersampling of larger extraterrestrial particles because of low sample time-area products may in some cases be a problem (see discussion in Peucker-Ehrenbrink and Ravizza 2000). The perhaps best value for the total accreted amount of extraterrestrial matter on the Earth's surface is 30,000 ($\pm 15,000$) metric tons per year (Peucker-Ehrenbrink and Ravizza 2000). This is based on Os isotopes in deep-sea sediments and agrees with the estimate of 30,000 ($\pm 10,000$) tons for the flux measured by the LDEF satellite at altitudes of 330 to 480 km (Love and Brownlee 1993; revised estimate in Engrand and Maurette 1998). Desert meteorite search programs show that the total mass flux to the Earth's surface for meteorites over the 10 g to 1 kg interval is between 2.9 and 7.3 tons per year, an estimate that agrees well with estimates from sky-watch networks (Bland et al. 1996; Halliday 2001). The large difference between the total and the meteorite-sized flux illustrates that the bulk of the extraterrestrial matter reaches the Earth surface as very small dust particles or micrometeorites ($<300 \mu\text{m}$), or occasionally as very large impactors ($>1\text{--}2 \text{ km}$).

Extraterrestrial chromite grains dispersed in condensed sediments is a new proxy for the influx of extraterrestrial detritus to Earth and has previously only been applied for mid-Ordovician condensed limestone (Schmitz et al. 2003; Schmitz and Haggström 2006). The average EC content of the Gubbio limestone, $0.029 \text{ grain kg}^{-1}$, is similar to that of mid-Ordovician limestone, ($0.013 \text{ grain kg}^{-1}$ for 379 kg) which formed before the disruption of the L chondrite parent body. Based on the Gubbio data, the average influx of EC to Earth is calculated to be $\sim 0.26 \text{ EC grain m}^{-2} \text{ kyr}^{-1}$, using average sedimentation rates of 2.5 mm kyr^{-1} for the Paleocene and 5 mm kyr^{-1} for the three Cretaceous samples studied (Arthur and Fischer 1977; Mukhopadhyay et al. 2001b) and a limestone density of 2.7 g cm^{-3} . For the mid-Ordovician, before the asteroid breakup event, the estimated flux is $\sim 0.09 \text{ EC grains m}^{-2} \text{ kyr}^{-1}$ but increases dramatically to an average of $\sim 13 \text{ EC grains m}^{-2} \text{ kyr}^{-1}$ after the event (Schmitz and Haggström 2006). We use a sedimentation rate of 2.5 mm kyr^{-1} for the Ordovician limestone, but this number contains an uncertainty of up to a factor of two. Furthermore, single EC grains can be readily overlooked during the chromite search and coincidental recovery of relict chromite from larger meteorite fragments, such as probably in sample +2.10 m at Gubbio, also adds statistical uncertainty to the flux estimates. We conclude that the normal flux of EC to Earth lies at $\sim 0.1\text{--}0.3 \text{ grain m}^{-2} \text{ kyr}^{-1}$.

The flux of EC grains ($0.26 \text{ grain m}^{-2} \text{ kyr}^{-1}$) reconstructed at Gubbio corresponds to $4.7 \times 10^{-4} \text{ g}$ of ordinary chondritic, unmelted debris $\text{m}^{-2} \text{ kyr}^{-1}$. For this estimate, we assume that the average EC grain is a cube with

a diameter of 100 μm ; we use a chromite density of 4.6 g cm^{-3} and an average chromite content for ordinary chondrites of 0.25 wt% (Keil 1962). Many of the EC grains have irregular shapes, but the use of a 100 μm cube in our calculation is reasonable, considering that all grains are $>63 \mu\text{m}$ in at least one dimension and most of the grains have maximum length axes $<150 \mu\text{m}$. The fact that we only search for chromite grains larger than 63 μm introduces some uncertainty, but for a gross, order-of-magnitude estimate the effect is negligible. Extrapolation to the entire surface of the Earth yields a global flux of ~ 200 tons of meteoritic matter per year. This is substantially less than the flux of $\sim 30,000$ tons based on Os isotopes in deep-sea sediments (Peucker-Ehrenbrink and Ravizza 2000), but significantly more than the 2.9–7.3 tons of meteorites in the 10 g to 1 kg mass spectrum according to desert meteorite abundances (Bland et al. 1996). Based on cosmic spherules from the South Pole Water Well, Taylor and Lever (2001) estimated an annual flux of 2700 ± 1400 tons of micrometeorites in the size range 50–700 μm ; however, 85% of the spherules were totally melted. Our mass estimate clearly appears realistic in view of these numbers and considering that our EC grains originate predominantly from decomposed unmelted meteorite fragments in the size range of 0.1 mm to a few centimeters (Schmitz et al. 2003; Heck et al., Forthcoming). The full range of the size spectrum and the significance of different size fractions within this spectrum are difficult to assess. Some micrometeorites may have been only marginally larger than the chromite grain itself. At the other end of the mass spectrum, chromite grains were spread over the sea floor by currents and bioturbation from decomposed kilogram-sized meteorites. Some of the chromite grains may derive from bodies 1–100 m in diameter, or even larger, which disrupted in the atmosphere or water column, resulting in micrometeorite and meteorite strewn fields on the sea floor (Lal and Jull 2002; Klekociuk et al. 2005).

Because of atmospheric disintegration of larger meteoroids, the size distribution of objects above the atmosphere is substantially different from that of the material reaching the Earth's surface (Lal and Jull 2002). In order to understand the processes that regulate the extraterrestrial flux through the atmosphere, it is necessary to understand the distribution of extraterrestrial matter over the different size fractions reaching Earth's surface. In flux estimates, the EC distribution in sediments appears to be an important complement to, for example, chemical studies of deep-sea sediments that primarily sample the most fine-grained and/or vaporized and recondensed fractions of the extraterrestrial flux (Lal and Jull 2003).

We stress that the flux of unmelted ordinary chondritic matter of 200 tons per year is a first-order estimate that may be refined as larger samples of limestone will be studied in the future. There are some potential methodological problems that have been considered. With the type of well-preserved

limestone like at Gubbio or in the mid-Ordovician at Kinnekulle in Sweden, there is good reason to believe that no EC grains have been destroyed over time by diagenesis or weathering (Schmitz and Häggström 2006). The EC grains found only show evidence of minor alteration, such as replacement of a few percent Fe by Zn. We do not find EC grains with, e.g., zoning that characterizes grains from more aggressive diagenetic environments, like in the Lockne impact crater where EC grains have been affected by hydrothermal activity (Alwmark and Schmitz 2007). Without grains representing intermediate states of alteration, it is not likely that grains were lost entirely because of diagenesis. Redistribution of EC grains by sea-floor currents may bias the EC grain distribution, but with large samples from several stratigraphic levels, the effect of this process will be smoothed out. Tests with individual chromite grains leached in HF and HCl show that the acids do not destroy the grains, but occasionally grains with fragile structure can break into smaller than 63 μm fragments during the wet sieving. Tests with samples "spiked" with EC grains show that this affects at the most a few percent of all grains.

CRETACEOUS-TERTIARY BOUNDARY EVENT

Chemical and isotopic extraterrestrial signatures at the K-T boundary worldwide and coeval species mass extinctions are best explained by the impact of a major extraterrestrial body ($\sim 10 \text{ km}$ in diameter) at the Yucatán Peninsula, Mexico (e.g., Alvarez et al. 1980; Hildebrand et al. 1991; Arenillas et al. 2006). The globally distributed K-T boundary clay containing iridium, shocked quartz, and spinel-rich microspherules represents compelling evidence for the impact event (Montanari and Koeberl 2000). Chromium isotopic analyses of the boundary clay (Shukolyukov and Lugmair 1998; Trinquier et al. 2006) and studies of a mm-sized weathered meteorite fragment from the K-T boundary in DSDP Hole 576 (Kyte 1998) suggest that the impactor was a carbonaceous chondrite, although a comet can not be ruled out. An important question is whether the K-T boundary impactor was a lone stray object or part of an asteroid or comet shower related to perturbations of solar system gravity (e.g., Hut et al. 1987). An enhanced flux of comets during a few million years can be produced by gravitational perturbations of the Oort Cloud, but the studies by Mukhopadhyay et al. (2001a, 2001b) at Gubbio indicate a near-constant flux of extraterrestrial ^3He during the K-T boundary event (Fig. 2), speaking against a comet shower. Varadi et al. (2003) suggest that the K-T impactor may instead have been related to perturbations in the asteroid belt $\sim 65 \text{ Myr}$ ago during chaotic transitions in the motions of the inner planets. Any such activity could have increased material transport in the inner solar system, resulting in enhanced accretion rates of cosmic material on Earth. High-resolution Ir analyses across the latest Maastrichtian and earliest

Paleocene strata at Gubbio, show that Ir is only enriched across a narrow interval close to the very K-T boundary, speaking against an extended period with substantially enhanced flux of extraterrestrial matter to Earth (Alvarez et al. 1990). Short vertical tails of enhanced Ir concentrations around the boundary clay most likely reflect redistribution of Ir during bioturbation and diagenesis. To add to this, our chromite searches in samples immediately adjacent to the K-T boundary clay could not find support for any significant increase in the flux of equilibrated ordinary chondritic micrometeorites or meteorites to Earth associated with the K-T event. The absence of $>63\text{ }\mu\text{m}$ chromite grains in beds immediately adjacent to the K-T boundary clay at Gubbio also speaks against an equilibrated ordinary chondritic impactor. Although we have not studied the centimeter-thick K-T boundary clay itself, chromite grains in adjacent beds can be expected if they occur in the boundary clay because of the effects of bioturbation and redistribution of material by currents. Alternatively, complete vaporization of an ordinary chondritic impactor during impact could have led to the absence of chromite. Our results are consistent with a carbonaceous chondritic impactor as suggested based on chromium isotopes. Carbonaceous chondrites only contain trace amounts of chromite, and grains larger than $63\text{ }\mu\text{m}$ are entirely absent or extremely rare (Fuchs et al. 1973; El Goresy 1976; Johnson and Prinz 1991; Rubin 1997), hence our data cannot rule out an increased flux of such material over an extended period.

CONCLUSIONS

The influx of extraterrestrial (ordinary chondritic) chromite ($>63\text{ }\mu\text{m}$) to Earth during the latest Maastrichtian and Paleocene has been established at a perfect testing ground, the condensed, pelagic limestones of the Bottaccione Gorge section at Gubbio, Italy. In 210 kg of limestone deposited at ~ 2.5 to 5 mm kyr^{-1} , we found a total of 6 EC grains, corresponding to an average of $0.03\text{ EC grain kg}^{-1}$. From well-constrained sedimentation rates, relying on detailed stratigraphic records, we determine the average flux of EC grains to Earth to $\sim 0.26\text{ grain m}^{-2}\text{ kyr}^{-1}$. This corresponds to a total influx of ~ 200 tons of extraterrestrial matter to Earth per year, which is realistic considering that the EC grains only record the flux of unmelted ordinary chondritic matter in the size range from $\sim 0.1\text{ mm}$ up to a few centimeters in diameter. The distribution of EC grains in sediments can give information on the extent to which extraterrestrial matter survives the passage through the atmosphere.

The results in this study confirm the extraordinary circumstances recorded in mid-Ordovician ($\sim 470\text{ Ma}$) limestone at Kinnekulle, Sweden, where an increase in EC grains of two orders of magnitude (from ~ 0.013 to $1\text{--}3\text{ EC grains kg}^{-1}$ limestone) coincides with the coeval disruption of the L chondrite parent body in the asteroid belt.

Large limestone samples collected at Gubbio in the immediate stratigraphic vicinity of the 1 cm thick K-T boundary clay contained no EC grains. This shows that there was no increase in the flux of ordinary chondritic matter to Earth during the K-T boundary transition.

The finding of 6 ordinary chondritic and 1 probable pallasitic chromite grain in 210 kg of condensed limestone from Gubbio shows that our approach has promise for retrieving general knowledge about past variations in the meteorite flux to Earth, particularly if methods could be developed for automatic preparation of limestone samples in the size of $100\text{--}200\text{ kg}$.

Acknowledgments—We are grateful to C. Alwmark for support at different occasions and to T. Häggström and J. Mellgren for field assistance at Gubbio. This study was funded by the Swedish Research Council and the Crafoord Foundation.

Editorial Handling—Dr. Timothy Swindle

REFERENCES

- Alvarez L. W., Alvarez W., Asaro F., and Michel H. V. 1980. Extraterrestrial cause for the Cretaceous-Tertiary extinction—Experimental results and theoretical interpretation. *Science* 208: 1095–1108.
- Alvarez W., Arthur M. A., Fischer A. G., Lowrie W., Napoleone G., Premoli Silva I., and Roggenthin W. M. 1977. Upper Cretaceous-Paleocene magnetic stratigraphy at Gubbio, Italy: V. Type section for the Late Cretaceous-Paleocene geomagnetic reversal time scale. *Geological Society of America Bulletin* 88: 383–389.
- Alvarez W., Asaro F., and Montanari A. 1990. Iridium profile for 10 million years across the Cretaceous-Tertiary boundary at Gubbio (Italy). *Science* 250:1700–1702.
- Alwmark C. and Schmitz B. 2007. Extraterrestrial chromite in the resurge deposits of the early Late Ordovician Lockne crater, central Sweden. *Earth and Planetary Science Letters* 253:291–303.
- Arenillas I., Arz J. A., Grajales-Nishimura J. M., Murillo-Muñeton G., Alvarez W., Camargo-Zanoguera A., Molina E., and Rosales-Domínguez C. 2006. Chicxulub impact event is Cretaceous/Paleogene boundary in age: New micropaleontological evidence. *Earth and Planetary Science Letters* 249:241–257.
- Arthur M. A. and Fischer A. G. 1977. Upper Cretaceous-Paleocene magnetic stratigraphy at Gubbio, Italy: I. Lithostratigraphy and sedimentology. *Geological Society of America Bulletin* 88:367–371.
- Barnes S. J. and Roeder P. L. 2001. The range of spinel compositions in terrestrial mafic and ultramafic rocks. *Journal of Petrology* 42: 2279–2302.
- Bell K., Castorina F., Lavecchia G., Rosatelli G., and Stoppa F. 2004. Is there a mantle plume below Italy? *Eos* 85:541; 545–547.
- Bevan A. W. R., Bland P. A., and Jull A. J. T. 1998. Meteorite flux on the Nullarbor region, Australia. In *Meteorites: Flux with time and impact effects*, edited by Grady M. M., Hutchison R., McCall G. J., and Rothery D. A. Geological Society London Special Publication #140. London: Geological Society of London. pp. 59–73.
- Bland P. A. 2001. Quantification of meteorite infall rates from

- accumulations in deserts, and meteorite accumulations on Mars. In *Accretion of extraterrestrial matter throughout Earth's history*, edited by Peucker-Ehrenbrink B. and Schmitz B. New York: Kluwer Academic/Plenum Publishers. pp. 267–303.
- Bland P. A., Smith T. B., Jull A. J. T., Berry F. J., Bevan A. W. R., Cloudt S., and Pillinger C. T. 1996. The flux of meteorites to the Earth over the last 50,000 years. *Monthly Notices of the Royal Astronomical Society* 283:551–565.
- Bridges J. C., Schmitz B., Hutchison R., Greenwood R. C., Tassinari M., and Franchi I. A. 2007. Petrographic classification of mid-Ordovician fossil meteorites from Sweden. *Meteoritics & Planetary Science* 43:1781–1789.
- Bunch T. E. and Keil K. 1971. Chromite and ilmenite in non-chondritic meteorites. *American Mineralogist* 56:146–157.
- Bunch T. E., Keil K., and Snetsinger K. G. 1967. Chromite composition in relation to chemistry and texture of ordinary chondrites. *Geochimica et Cosmochimica Acta* 31:1569–1582.
- Cresta S., Monechi S., and Parisi G. 1989. Mesozoic-Cenozoic stratigraphy in the Umbria-Marche area. *Memorie Descrittive della Carta Geologica d'Italia* 39:1–185.
- Dalai T. K. and Ravizza G. 2006. Evaluation of osmium isotopes and iridium as paleoflux tracers in pelagic carbonates. *Geochimica et Cosmochimica Acta* 70:3928–3942.
- Ebel D. S. and Grossman L. 2005. Spinel-bearing spherules condensed from the Chicxulub impact-vapor plume. *Geology* 33: 293–296.
- El Goresy A. 1976. Opaque oxide minerals in meteorites. In *Oxide minerals*, edited by Rumble D. III. Reviews in Mineralogy, vol. 3. Washington, D.C.: Mineralogical Society of America. pp. EG47–EG72.
- Engrand C. and Maurette M. 1998. Carbonaceous micrometeorites from Antarctica. *Meteoritics & Planetary Science* 33:565–580.
- Farley K. A., Love S. G., and Patterson D. B. 1997. Atmospheric entry heating and helium retentivity of interplanetary dust particles. *Geochimica et Cosmochimica Acta* 61:2309–2316.
- Farley K. A., Montanari A., Shoemaker E. M., and Shoemaker C. S. 1998. Geochemical evidence for a comet shower in the late Eocene. *Science* 280:1250–1253.
- Farley K. A., Vokrouhlicky D., Bottke W. F., and Nesvorny D. 2006. A late Miocene dust shower from the breakup of an asteroid in the main belt. *Nature* 439:295–297.
- Fuchs L. H., Olsen E., and Jensen K. J. 1973. Mineralogy, mineral-chemistry, and composition of the Murchison (C2) meteorite. *Smithsonian Contributions to the Earth Sciences* 10:1–39.
- Gabrielli P., Barbante C., Plane J. M. C., Varga A., Hong S., Cozzi G., Gaspari V., Planchon F. A. M., Cairns W., Ferrari C., Crutzen P., Cescon P., and Boutron C. F. 2004. Meteoric smoke fallout over the Holocene epoch revealed by iridium and platinum in Greenland ice. *Nature* 432:1011–1014.
- Greenwood R. C., Schmitz B., Bridges J. B., Hutchison R., and Franchi I. A. 2007. Disruption of the L-chondrite parent body: New oxygen isotope evidence from Ordovician relict chromite grains. *Earth and Planetary Science Letters* 262:204–213.
- Halliday I. 2001. The present-day flux of meteorites to the Earth. In *Accretion of extraterrestrial matter throughout Earth's history*, edited by Peucker-Ehrenbrink B. and Schmitz B. New York: Kluwer Academic/Plenum Publishers. pp. 305–318.
- Heck P. R., Schmitz B., Baur H., Halliday A. N., and Wieler R. 2004. Fast delivery of meteorites to Earth after a major asteroid collision 480 Myr. *Nature* 430:323–325.
- Heck P. R., Schmitz B., Baur H., and Wieler R. Forthcoming. Noble gases in fossil micrometeorites and meteorites from 470 Myr old sediments from southern Sweden and new evidence for the L chondrite parent body breakup event. *Meteoritics & Planetary Science* 43.
- Hildebrand A. R., Penfield G. T., Kring D. A., Pilkington M., Camargo A., Jacobsen S. B., and Boynton W. V. 1991. Chicxulub crater: A possible Cretaceous-Tertiary boundary impact crater on the Yucatán Peninsula, Mexico. *Geology* 19: 867–871.
- Hills J. G. 1981. Comet showers and the steady-state infall of comets from the Oort cloud. *The Astronomical Journal* 86: 1730–1740.
- Hut P., Alvarez W., Elder W. P., Hansen T., Kauffman E. G., Keller G., Shoemaker E. M., and Weissman P. R. 1987. Comet showers as cause of mass extinctions and group differences.
- Jarosewich E., Nelen J. A., and Norberg J. A. 1980. Reference samples for electron microprobe analysis. *Geostandards Newsletter* 4:43–47.
- Johnson C. A. and Prinz M. 1991. Chromite and olivine in type II chondrules in carbonaceous and ordinary chondrites: Implications for thermal histories and group differences. *Geochimica et Cosmochimica Acta* 55:893–904.
- Karner D. B., Levine J., Muller R. A., Asaro F., Ram M., and Stolz M. R. 2003. Extraterrestrial accretion from the GISP2 ice core. *Geochimica et Cosmochimica Acta* 67:751–763.
- Keil K. 1962. On the phase composition of meteorites. *Journal of Geophysical Research* 67:4055–4061.
- Klekociuk A. R., Brown P. G., Pack D. W., ReVelle D. O., Edwards W. N., Spalding R. E., Tagliaferri E., Yoo B. B., and Zagari J. 2005. Meteoritic dust from the atmospheric disintegration of a large meteoroid. *Nature* 436:1132–1135.
- Kuhnt W. 1990. Agglutinated foraminifera of western Mediterranean Upper Cretaceous pelagic limestone (Umbrian Apennines, Italy, and Betic Cordillera, southern Spain). *Micropaleontology* 36: 297–330.
- Kyte F. T. 1998. A meteorite from the Cretaceous/Tertiary boundary. *Nature* 396:237–239.
- Kyte F. T. and Smit J. 1986. Regional variations in spinel compositions: An important key to the Cretaceous/Tertiary event. *Geology* 14:485–487.
- Lal D. and Jull A. J. T. 2002. Atmospheric cosmic dust fluxes in the size range 10^{-4} to 10 centimeters. *The Astrophysical Journal* 576: 1090–1097.
- Lal D. and Jull A. J. T. 2003. Extraterrestrial influx rates of cosmogenic isotopes and platinum group elements: Realizable geochemical effects. *Geochimica et Cosmochimica Acta* 67: 4925–4933.
- Love S. G. and Brownlee D. E. 1993. A direct measurement of the terrestrial mass accretion rate of cosmic dust. *Science* 262:550–553.
- Lowrie W., Alvarez W., and Asaro F. 1990. The origin of the White Beds below the Cretaceous-Tertiary boundary in the Gubbio section, Italy. *Earth and Planetary Science Letters* 98: 303–312.
- Matese J. J., Whitman P. G., Innanen K. A., and Valtonen M. J. 1995. Periodic modulation of the Oort cloud comet flux by the adiabatically changing galactic tide. *Icarus* 116:225–268.
- Mathews J. D., Janches D., Meisel D. D., and Zhou Q. H. 2001. The micrometeoroid mass flux into the upper atmosphere: Arecibo results and a comparison with prior estimates. *Geophysical Research Letters* 28:1929–1932.
- Monechi S. and Thierstein H. R. 1985. Late Cretaceous-Eocene nannofossil and magnetostratigraphic correlations near Gubbio, Italy. *Marine Micropaleontology* 9:419–440.
- Montanari A. 1991. Authigenesis of impact spheroids in the K/T boundary clay from Italy: New constraints for high-resolution stratigraphy of terminal Cretaceous events. *Journal of Sedimentary Petrology* 61:315–339.
- Montanari A. and Koeberl C. 2000. *Impact stratigraphy—The Italian*

- record. Lecture Notes in Earth Sciences, vol. 93. Heidelberg: Springer. 364 p.
- Mukhopadhyay S., Farley K. A., and Montanari A. 2001a. A short duration of the Cretaceous-Tertiary boundary event: Evidence from extraterrestrial ^3He . *Science* 291:1952–1955.
- Mukhopadhyay S., Farley K. A., and Montanari A. 2001b. A 35 Myr record of helium in pelagic limestone from Italy: Implications for interplanetary dust accretion from the early Maastrichtian to the middle Eocene. *Geochimica et Cosmochimica Acta* 65:653–669.
- Nesvorný D., Vokrouhlický D., Bottke W. F., and Sykes M. 2006. Physical properties of asteroid dust bands and their sources. *Icarus* 181:107–144.
- Nyström J. O., Lindström M., and Wickman F. E. 1988. Discovery of a second Ordovician meteorite using chromite as a tracer. *Nature* 336:572–574.
- Peucker-Ehrenbrink B. 2001. Iridium and osmium as tracers of extraterrestrial matter in marine sediments. In *Accretion of extraterrestrial matter throughout Earth's history*, edited by Peucker-Ehrenbrink B. and Schmitz B. New York: Kluwer Academic/Plenum Publishers. pp. 163–178.
- Peucker-Ehrenbrink B. and Ravizza G. 2000. The effects of sampling artifacts on cosmic dust flux estimates: A reevaluation of nonvolatile tracers (Os, Ir). *Geochimica et Cosmochimica Acta* 64:1965–1970.
- Premoli Silva I. 1977. Upper Cretaceous-Paleocene magnetic stratigraphy at Gubbio, Italy; II. Biostratigraphy. *Geological Society of America Bulletin* 88:371–374.
- Premoli Silva I. and Sliter W. V. 1995. Cretaceous planktonic foraminiferal biostratigraphy and evolutionary trends from the Bottaccione section, Gubbio, Italy. *Palaeontographia Italica* 82: 1–89.
- Ramdohr P. 1973. *The opaque minerals in stony meteorites*. Amsterdam: Elsevier. 245 p.
- Robin E., Bonté Ph., Froget L., Jéhanno C., and Rocchia R. 1992. Formation of spinels in cosmic objects during atmospheric entry: A clue to the Cretaceous-Tertiary boundary event. *Earth and Planetary Science Letters* 108:181–190.
- Roeder P. L. 1994. Chromite: From the fiery rain of chondrules to the Kilauea Iki lava lake. *The Canadian Mineralogist* 32:729–746.
- Rubin A. E. 1997. Mineralogy of meteorite groups. *Meteoritics & Planetary Science* 32:231–247.
- Schmitz B. and Häggström T. 2006. Extraterrestrial chromite in Middle Ordovician marine limestone at Kinnekulle, southern Sweden—Traces of a major asteroid breakup event. *Meteoritics & Planetary Science* 41:455–466.
- Schmitz B., Lindström M., Asaro F., and Tassinari M. 1996. Geochemistry of meteorite-rich marine limestone strata and fossil meteorites from the lower Ordovician at Kinnekulle, Sweden. *Earth and Planetary Science Letters* 145:31–48.
- Schmitz B., Tassinari M., and Peucker-Ehrenbrink B. 2001. A rain of ordinary chondritic meteorites in the early Ordovician. *Earth and Planetary Science Letters* 194:1–15.
- Schmitz B., Häggström T., and Tassinari M. 2003. Sediment-dispersed extraterrestrial chromite traces a major asteroid disruption event. *Science* 300:961–964.
- Shukolyukov A. and Lugmair G. W. 1998. Isotopic evidence for the Cretaceous-Tertiary impactor and its type. *Science* 282:927–929.
- Smit J. 1982. Extinction and evolution of planktonic foraminifera after a major impact at the Cretaceous/Tertiary boundary. GSA Special Paper #190. Boulder, Colorado: Geological Society of America. pp. 329–352.
- Smit J. and Kyte F. T. 1984. Siderophile-rich magnetic spheroids from the Cretaceous-Tertiary boundary in Umbria, Italy. *Nature* 310:403–405.
- Taylor S. and Lever J. H. 2001. Seeking unbiased collections of modern and ancient micrometeorites. In *Accretion of extraterrestrial matter throughout Earth's history*, edited by Peucker-Ehrenbrink B. and Schmitz B. New York: Kluwer Academic/Plenum Publishers. pp. 205–219.
- Thorslund P., Wickman F. E., and Nyström J. O. 1984. The Ordovician chondrite from Brunflo, central Sweden. I. General description and primary minerals. *Lithos* 17:87–100.
- Toppani A. and Libourel G. 2003. Factors controlling composition of cosmic spinels: Application to atmospheric entry conditions of meteoritic materials. *Geochimica et Cosmochimica Acta* 67: 4621–4638.
- Trinquier A., Birck J.-L., and Allègre C. J. 2006. The nature of the K-T impactor. A ^{54}Cr reappraisal. *Earth and Planetary Science Letters* 241:780–788.
- Varadi F., Runnegar B., and Ghil M. 2003. Successive refinements in long-term integrations of planetary orbits. *The Astrophysical Journal* 592:620–630.
- Wlotzka F. 2005. Cr spinel and chromite as petrogenetic indicators in ordinary chondrites: Equilibration temperatures of petrologic types 3.7 to 6. *Meteoritics & Planetary Science* 40: 1673–1702.
- Zappalà V., Cellino A., Gladman B. J., Manley S., and Migliorini F. 1998. Asteroid showers on Earth after family breakup events. *Icarus* 134:176–179.